# Quantabio

Product Information		
PerfeC⊤a <sup>®</sup> qPCR SuperMix, ROX™		
Part Number	95051-100	
Number of Reactions	100 reactions	
Reaction Size	50 μL	
Storage Temperature	-25ºC to -15ºC	
Lot Number	66147777	
Reference Number	091619	
Expiration Date	09/30/2022	

### Product Description:

PerfeCta qPCR SuperMix ROX is a 2X concentrated, readyto-use reaction cocktail that contains all components, except primers, probe(s), and template for real-time quantitative PCR systems. The proprietary buffer and stabilizers have been specifically optimized to deliver maximum PCR efficiency, sensitivity, and robust fluorescent signal with TaqMan® or TaqMan MGB probe chemistry. The enhanced specificity of this supermix suppresses crossreactivity between homologous sequences, improving detection and discrimination in SNP applications. A key component of this supermix is AccuStart<sup>™</sup> Taq DNA polymerase, which contains monoclonal antibodies that bind to the polymerase and keep it inactive prior to the initial PCR denaturation step. Upon heat activation (2 minutes at 95°C), the antibodies denature irreversibly, releasing fully active, unmodified Taq DNA polymerase. This enables specific and efficient primer extension with the convenience of room temperature reaction assembly.

## <u>Component Part Numbers:</u> 84010 PerfeCta qPCR SuperMix, ROX 1.25mL

Product Specifications			
95051			
Assay	qPCR ß-actin Plasmid DNA Functional Assay	DNase	RNase
Result	Pass	Pass	Pass

## **Quality Control Analysis and Specifications:**

### Nuclease Assay:

**DNase:** DNase activity must be below the detectable limits of 100 pg DNase I equivalent as assayed using a fluorogenic substrate following a 1 hour incubation at 37°C with each kit component at 1X concentration.

**RNase:** RNase activity must be below the detectable limits of 1 pg RNase A equivalent as assayed using a fluorogenic substrate following a 1 hour incubation at 37°C with each kit component at 1X concentration.

**qPCR** *B***-actin Plasmid DNA Functional Assay:** Detection of  $\beta$ -actin from 10 copies to 1 x 10<sup>7</sup> copies. Correlation of determination (R<sup>2</sup>)  $\geq$  0.990 from Ct standard curve analysis. Slope from Ct standard curve analysis between -3.20 and -3.70.

### Limitations of Use

Quantabio and Ultraplex are registered trademarks of QIAGEN Beverly, Inc. Quanta Biosciences, qScript, Geltrack, ToughMix, PerfeCta, and Fastmix are registered trademarks of Quanta BioSciences Inc. Extracta, AccuStart, AccuMelt, and Accuvue are trademarks of Quanta BioSciences Inc. Applied Biosystems, StepOne, StepOnePlus and ROX are trademarks of Thermo Fisher Scientific and or its subsidiaries. Please contact QIAGEN-Beverly for more information.

This product was developed, manufactured, and sold for *in vitro* use only. The product is not suitable for administration to humans or animals. SDS sheets relevant to this product are available upon request. 100 Cummings Center, Suite 407J, Beverly, MA 01915 • Ph (888) 927-7027 • Fax (978) 867-5724 • <u>www.QuantaBio.com</u> FMWI016.2 Rev 01